#### 7/ 18

### IN THE SPECIFICATION:

### Please insert the following paragraph on Page 8, line 28, as follows:

-- From a mathematical standpoint, there are infinite numbers of functions (f(x,y) that can approximately map the two spaces 12 and 14, as shown in the literature. This invention proposes a simple function that follows the predictions, as set forth herein.--

## Please insert the following paragraph on Page 9, line 1 (below the graph), as follows:

--Graphically presented is as follows consider two points in the panel before pressing  $M(x_1,y_1)$  and  $N(x_2,y_2)$ , after pressing the two points move to new locations  $M'(x'_1,y'_1)$  and  $N'(x'_2,y_{22})$ .

Note that after pressing, the angle (MOH) =  $\arctan(y_1/x_1)$  changes to new value  $\arctan(y'_1/x'_1)$ .

The line MN translates, scales (stretches or compresses), and rotates an angle  $\alpha$  to a new location M'N'. Note that in this example, the line MN is stretched due to the increasing in length.--

# Please revise the following paragraph beginning on Page 10, line 6, to read as follows:

5-17-06 Py

--It is possible then to easily determine the coefficients  $A_x$ ,  $A_y$ ,  $B_x$ ,  $B_y$ ,  $C_x$ ,  $C_y$ ,  $D_x$  and  $D_y$  as follows. Using the same origin, for i=1, 2, 3 and 4, let  $(x_i, y_i)$  and  $(x'_i, y'_i)$  be the coordinates of four known points before pressing and after pressing, respectively. Writing equations (3) and (4) for x- and y- directions, wherein:--

#### 8/ 18

Please insert the following paragraph on Page 10, line 20, as follows:

--Solve the above eight independent equations for eight unknowns A<sub>x</sub>, A<sub>y</sub>, B<sub>x</sub>, B<sub>y</sub>, C<sub>z</sub>, C<sub>y</sub>, D<sub>z</sub> and D<sub>y</sub>.

Then substitute them to equations (3) and (4).--

Please revise the following paragraph beginning on Page 14, line 6, to read as follows:

--Computing the angle between the line P'Q' and R'S'. Equation of the line passing P'Q' is  $y=-2.2702(10)^{-7} x + 9.9955$ , and equation of the line passing R'S' is  $y=1.2432(10)^{-3} x + 322.78$ . Take two vectors:  $\{1, -2.2702(a0)-7\}^T$  points along P'Q' and  $\{1, -1.2432(10)-3\}^T$  points along R'S'. Then the angle between these two vectors is 0.001243 radians.--